

Original

## Midazolam inhibits IgE production in mice via suppression of class switch recombination

Hiroto Kusama<sup>1)</sup>, Ryoki Kobayashi<sup>2)</sup>, and Tomoko Kurita-Ochiai<sup>2)</sup>

<sup>1)</sup>Department of Anesthesiology, Nihon University School of Dentistry at Matsudo, Matsudo, Japan

<sup>2)</sup>Department of Oral Immunology, Nihon University School of Dentistry at Matsudo, Matsudo, Japan

(Received October 18, 2013; Accepted January 29, 2014)

**Abstract:** Anaphylactic shock is characterized by increased capillary permeability and a decline in blood pressure due to excessive production of IgE. Midazolam (MDZ) is reported to have immunomodulatory properties. However, little is known about the effect of MDZ on the production of IgE antibody. We examined whether MDZ can suppress antigen-specific and total IgE production followed by IgE class switch recombination (CSR). MDZ was administered intraperitoneally to mice prior to ovalbumin (OVA) plus native cholera toxin (nCT) immunization. Serum OVA-specific and total IgE responses, and surface IgE-positive B cells were analyzed by ELISA and flow cytometry. Furthermore, expression levels of CSR-associated molecules such as germ-line transcript  $\epsilon$  ( $\epsilon$ GLT), germ-circle transcript  $\epsilon$  ( $\epsilon$ CT), AID, and Id2 in the spleen were compared. The levels of interferon-gamma (IFN- $\gamma$ ) and interleukin (IL)-4 mRNA and protein were also examined in the spleen and serum. MDZ significantly suppressed OVA-specific and total IgE levels in plasma and surface IgE-positive B cells in the spleen. Moreover, MDZ-treated mice had significantly reduced levels of  $\epsilon$ GLT and  $\epsilon$ CT. Furthermore, although the levels of IFN- $\gamma$  mRNA and protein were significantly elevated, those of IL-4 were reduced in MDZ-treated mice. Therefore, MDZ may be an important modulator of allergic responses through its ability to downregulate IgE production. (J Oral Sci 56, 77-83, 2014)

Keywords: midazolam; IgE; OVA; spleen B cell.

### Introduction

Immunoglobulin E (IgE) plays a central role in the pathogenesis of many allergic diseases, such as allergic asthma, allergic rhinitis, and atopic dermatitis. Furthermore, immediate hypersensitivity reactions to anesthetics and associated agents used during the perioperative period have been reported with increasing frequency in most developed countries (1). In sensitized individuals, both total and allergen-specific IgE antibodies (Abs) are produced at high levels and bind to high-affinity Fc $\epsilon$ Rs on mast cells and basophils surfaces, leading to the release of preformed and newly synthesized mediators that initiate immunologic cascades and inflammatory reactions. Therefore, inhibition of IgE production is an ideal strategy for amelioration of allergic disease. Until now, no therapies capable of suppressing IgE production have been clinically available, although neutralization of pre-existing IgE has been applied to patients with severe allergic asthma (2). Although IgE has a relatively short half-life in plasma, mechanisms that tightly control IgE class switch recombination (CSR) are thought to account for the low levels of IgE normally observed in plasma (3-5). CSR replaces the heavy chain constant region C $\mu$  gene with a targeted C $\epsilon$  gene by recombination of a switch region  $\mu$  (S $\mu$ ) with a S $\epsilon$  region present in the respective targeted CH gene. Consequently, CSR allows the expression of an Ig that has the same antigen specificity with a secondary heavy chain isotype (IgG, IgA, or IgE) that can exhibit different effector functions. Regulation of CSR in B cells is coordinated with the germ line transcription (GLT) of CH genes, as well as the induction of activation-induced cytidine deaminase (AID) expression (6).

Correspondence to Dr. Tomoko Kurita-Ochiai, Department of Oral Immunology, Nihon University School of Dentistry at Matsudo, 2-870-1 Sakaecho-nishi, Matsudo, Chiba 271-8587, Japan  
Fax: +81-47-360-9601 E-mail: ochiai.tomoko@nihon-u.ac.jp

doi.org/10.2334/josnusd.56.77  
DN/JST.JSTAGE/josnusd/56.77

Midazolam (MDZ), a benzodiazepine (BZD), is an intravenously administered anesthetic used for premedication, induction and maintenance of general anesthesia, as well as for sedation in the treatment of nervous dental patients. There are two types of BZD receptors, central-type BZD receptors (CBRs) and peripheral-type BZD receptors (PBRs) (7,8). CBRs are expressed mainly in the central nervous system (7), while PBRs are detected in many peripheral tissues, cells and organs, such as the kidney, endocrine organs and monocytes (7,8). PBRs are reportedly involved in regulation of cellular proliferation, immunomodulation, steroidogenesis, and oxidative processes, as well as programmed cell death (7,8). MDZ has also been shown to inhibit IL-6 mRNA expression in human peripheral blood mononuclear cells (9), and to suppress lipopolysaccharide (LPS)-induced release of nitric oxide and TNF- $\alpha$  from rat microglia via PBRs (10). These results led us to speculate that MDZ might modulate immune system function. However, the exact mechanism responsible for the effects of MDZ on IgE production remains to be fully elucidated.

In the present study, therefore, we examined the effect of i.p.-administrated MDZ on total and antigen-specific levels of IgE as well as IgE CSR in serum and splenocytes from ovalbumin (OVA) plus cholera toxin (CT)-immunized BALB/c mice. Furthermore, we investigated the immunomodulatory activities of MDZ on induced interferon-gamma (IFN- $\gamma$ ) and interleukin-4 (IL-4) production and expression of their mRNAs in serum and splenocytes from OVA-immunized BALB/c mice.

## Materials and Methods

### Mice

Female BALB/c Cr Slc mice were purchased from Sankyo Laboratories (Tokyo, Japan) and maintained under specific pathogen-free conditions at an experimental facility of Nihon University School of Dentistry at Matsudo, Chiba, Japan. Mice were 8 to 12 weeks old when used for the experiments. All food and water were sterile. All animals were maintained and used in accordance with the guidelines for the care and use of laboratory animals of Nihon University School of Dentistry at Matsudo (AP13MD007).

### Optimization of MDZ dose

In humans, the loading dose of MDZ is 0.15 mg/kg, followed by continuous infusion of 1–7  $\mu$ g/kg/min. The maximum dose is 6 to 10 mg/kg (11,12). Therefore, each mouse (five mice/group) was administered different intraperitoneal (i.p.) doses (1 mg, 3 mg, or 10 mg/kg)

of MDZ in 100  $\mu$ L phosphate-buffered saline (PBS) in order to establish the minimum effective dose of MDZ. Mice were monitored daily for staggering, reduction of physical activity, sedation, and loss of the righting reflex.

### Immunization and sample collection

BALB/c mice (five mice/group) were administered MDZ (3 mg/kg) or PBS alone via i.p. injection and then immunized intraperitoneally with 1 mg of ovalbumin (OVA, fraction V; Sigma-Aldrich, St. Louis, MO, USA) plus 1  $\mu$ g of native cholera toxin (nCT) in PBS on days 0, 7, and 14. On day 21, serum samples were collected and mononuclear cells were isolated from the spleens of both MDZ-treated and untreated mice that had been immunized with OVA and nCT.

### OVA-specific and cytokine-specific ELISA

The presence of OVA-specific IgE and total IgE Abs in serum was determined using an ELISA kit (Rebisu, Shibayagi, Shibukawa, Japan). Levels of cytokines in serum were measured using ELISA. A Quantikine mouse immunoassay kit (R&D Systems, Minneapolis, MN, USA) was used to detect IFN- $\gamma$  and IL-4 in the plasma of both MDZ-treated and control mice immunized with OVA plus nCT.

### Flow cytometry

To characterize the frequency of surface-IgE-positive (sIgE<sup>+</sup>) B cells,  $2 \times 10^5$  mononuclear cells were incubated with phycoerythrin (PE)-labeled anti-IgE Ab and allophycocyanin-tagged B220 (BioLegend, San Diego, CA, USA). Samples were then subjected to FACS analysis (BD Biosciences, San Jose, CA, USA).


### Quantitative real-time PCR analysis

Total RNA was purified from splenic cells using Trizol reagent (Invitrogen, Tokyo, Japan) in accordance with the manufacturer's protocol. Superscript II reverse transcriptase (Invitrogen, Carlsbad, CA, USA) was used to generate cDNA from purified total RNA obtained from each mouse. Quantitative real-time RT-PCR analyses were performed using a Thermal Cycler Dice real-time PCR system (Takara, Shiga, Japan) in accordance with the manufacturer's protocol. The initial denaturation step was carried out at 95°C for 10 min, followed by 40 cycles of 95°C for 15 s, 60°C for 25 s, and 60°C for 35 s. Each gene was tested in triplicate. Target RNA levels were normalized to that of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA. The levels of germ-line transcript  $\epsilon$  ( $\epsilon$ GLT), germ-circle transcript  $\epsilon$  ( $\epsilon$ CT), AID, Id2, IFN- $\gamma$ , IL-4, and GAPDH cDNA were determined

**Table 1** Optimization of MDZ dose

Midazolam	Stagger	Decrease in physical activity	Sedation	Loss of righting reflex
1 mg/kg	2/5	3/5	1/5	5/5
3 mg/kg	4/5	5/5	4/5	5/5
10 mg/kg	5/5	5/5	5/5	5/5

Various doses (1 mg, 3 mg, or 10 mg/kg/mouse) of MDZ were administered intraperitoneally to BALB/c mice (five mice per group). Ten minutes later, behavioral effects induced by MDZ were examined.

**Fig. 1** Serum IgE response.

BALB/c mice were pre-administered MDZ (3 mg/kg) or PBS alone via i.p. injection and then immunized with OVA plus nCT on days 0, 7, and 14 via i.p. injection. Serum samples were collected on day 21, and the OVA-specific and total IgE responses were examined by ELISA. Results are expressed as the mean  $\pm$  SD of five mice per group. \*\*\* $P < 0.001$  compared with control (OVA plus nCT + PBS).

by quantitative PCR using the following primer pairs:

$\epsilon$ GLT: 5'-GCACAGGGGGCAGAAGAT-3', 5'-CCA-GGGTCATGGAAGCAGTG-3'

$\epsilon$ CT: 5'-TTGGACTACTGGGGTCAAGG-3', 5'-CA-GTGCCCTTACAGGGCTTC-3'

AID: 5'-GGCTGAGGTTAGGGTTCATCTCAG-3', 5'-GAGGGAGTCAAGAAAGTCACGCTGGA-3'

Id2: 5'-ATCGTCTTGCCAGGTGTCGTTCT-3', 5'-AGCATCCCCAGAACAAGAAGGTG-3'

IFN- $\gamma$ : 5'-CGGCACAGTCATTGAAAGCCTA-3', 5'-GTTGCTGATGGCCTGATTGTC-3'

IL-4: 5'-CATCGGCATTTGAACGAGGTCA-3', 5'-CTTATCGATGAATCAGGCATCG-3'

GAPDH: 5'-TGTGTCCGTCGTGGATCTGA-3', 5'-TTGCTGTTGAAGTCGCAGGAG-3'

### Statistical analysis

Student's *t*-test (two-tailed) was used for all statistical analyses, and differences at  $P < 0.05$  were considered statistically significant.

## Results

### MDZ inhibits Ag-specific IgE responses in serum


We examined the optimal dose of i.p.-administered MDZ. Various doses (1 mg, 3 mg, and 10 mg/kg/mouse) of MDZ were administered i.p. to five mice in each group. Of the five mice that were given a 3 mg/kg MDZ, at least four demonstrated evidence of anesthesia, such as staggering, a reduction in physical activity, sedation, and/or loss of the righting reflex (Table 1). As 3 mg/kg MDZ appeared to be the optimal concentration for inducing a hypnotic-anesthetic state, this concentration was used in all subsequent experiments. To evaluate Ag-specific IgE responses, mice were given MDZ i.p. 30 min before being immunized with OVA plus nCT (also via the i.p. route). Serum was collected from each mouse 7 days after the third i.p. injection, after which IgE levels were determined using ELISA. As shown in Fig. 1, MDZ inhibited OVA-specific and total IgE production. This effect of MDZ on IgE production reflected a reduction of surface IgE expression on B cells, as confirmed by FACS analysis (Fig. 2).

### Suppression of IgE CSR in MDZ-treated mice

We examined the effects of MDZ on CSR-mediated regulatory genes in spleen cells of mice that had been immunized with OVA plus nCT. The levels of mRNA specific for  $\epsilon$ GLT,  $\epsilon$ CT, AID, and Id2 were determined using quantitative real-time PCR. Spleen cells from mice not treated with MDZ served as a positive control. The level of AID-specific mRNA was comparable in MDZ-treated and untreated mice (Fig. 3). In contrast, significant inhibition of  $\epsilon$ GLT (A) and  $\epsilon$ CT (B) mRNA production was observed in MDZ-treated mice relative to control mice. Interestingly, Id2, which suppresses IgE CSR, was increased in MDZ-treated mice. These data indicate that inhibition of IgE CSR by MDZ is caused by suppression of  $\epsilon$ GLT and  $\epsilon$ CT synthesis.


### Induction of IFN- $\gamma$ but not IL-4 expression in spleen cells

Since the number of surface IgE<sup>+</sup> B cells and the frequency of IgE CSR were lower in MDZ-treated mice than in



**Fig. 2** Frequency of surface IgE-positive B cells.

BALB/c mice were pre-administered MDZ or PBS and then immunized with OVA plus nCT or PBS alone, as described in the legend to Fig. 1. Splens were collected on day 21 and single-cell suspensions were prepared. The cells were stained with Alex Fluor 647- conjugated anti-mouse B220 and PE-conjugated anti-mouse IgE, and analyzed by flow cytometry on a FSC/B220 plot (A) or IgE/B220 plot (B). The B cell population (Gate R1) was further gated for B220 binding and forward scatter (FSC) properties (A) by selecting those that were stained with anti-IgE Abs. The results are representative of five experiments. Panel C shows the summarized results for five mice per group. Results are expressed as the mean  $\pm$  SD \*\*\* $P$  < 0.001 compared with control (OVA plus nCT + PBS).




**Fig. 3** Expression of class switch recombination-associated molecules.

BALB/c mice were pre-administered MDZ or PBS and then immunized with OVA plus nCT as described in the legend to Fig. 1. Splens were removed on day 21 and total RNA was isolated. The levels of  $\epsilon$ GLT (A),  $\epsilon$ CT (B), AID (C), and Id2 (D) mRNA transcripts were determined using quantitative RT-PCR. Relative mRNA levels were determined after normalization to the level of GAPDH mRNA. Values represent mean values  $\pm$  SD ( $n = 5$ ). \* $P$  < 0.05 compared with control (OVA plus nCT + PBS). \*\* $P$  < 0.005 compared with control (OVA plus nCT + PBS). \*\*\* $P$  < 0.001 compared with control (OVA plus nCT + PBS).

untreated controls, we speculated that some soluble factors might be involved in MDZ-mediated suppression of CSR. Therefore we assessed the production of the cytokines IFN- $\gamma$  and IL-4, which are known to regulate IgE CSR in activated B cells (13). After confirming that the level of OVA-specific IgE in serum was decreased in MDZ-treated mice on day 21 (7 days after the last immunization), spleen cells were collected and their levels of IFN- $\gamma$  and IL-4 mRNA were determined using real time-PCR. Significant induction of IFN- $\gamma$ -specific mRNA transcript production was detected in the spleen cells of MDZ-treated mice (Fig. 4). However, the level of IL-4-specific mRNA was significantly reduced in MDZ-treated mice relative to control mice. Moreover, in MDZ-treated mice, the altered levels of IFN- $\gamma$  and IL-4 cytokines in serum were correlated with mRNA transcript levels (Fig. 5).

## Discussion


In this study, we demonstrated that MDZ inhibits the production of OVA-specific IgE in serum as well as the production of surface IgE<sup>+</sup> B cells in the spleen, in concert with suppression of IgE CSR. Moreover, increased levels of IFN- $\gamma$  mRNA and decreased levels of IL-4 mRNA, with corresponding changes in the levels of the respective proteins were detected in serum and spleen cells from MDZ-treated mice. These data suggest that inhibition of Ag-specific IgE production by MDZ is associated with differential regulation of cytokines. Therefore we focused on MDZ-mediated immunomodulation of Th1/Th2 cell balance and suppression of IgE production. It now seems that Th1 and Th2 responses are strongly tied to IFN- $\gamma$



**Fig. 4** Expression of IFN- $\gamma$  and IL-4 mRNA.

BALB/c mice were pre-administered MDZ or PBS and then immunized with OVA plus nCT, as described in the legend to Fig. 1. Spleens were removed on day 21 and total RNA was isolated. The levels of IFN- $\gamma$  (A) and IL-4 (B) mRNA transcripts were determined using quantitative RT-PCR. Relative mRNA levels were determined after normalization to the level of GAPDH mRNA using real-time RT-PCR. Values represent the mean values  $\pm$  SD ( $n = 5$ ). \*\*\* $P < 0.001$  compared with control (OVA plus nCT + PBS).

and IL-4, respectively. We first examined the ability of MDZ to modulate the *in vivo* production of IFN- $\gamma$  and IL-4 in spleen cells from BALB/c mice immunized with OVA plus nCT in order to clarify the potential of MDZ to induce a Th1-skewed response. In this trial, MDZ administration effectively reduced the level of OVA-specific IgE in serum (Fig. 1). In parallel with suppression of IgE in serum, i.p. administration of MDZ resulted in increased IFN- $\gamma$  production concomitant with suppression of IL-4 production in serum (Fig. 5). The reduction in serum IgE could be explained by enhanced IFN- $\gamma$  production and suppressed IL-4 production, since IL-4 signaling is a prerequisite for IgE synthesis in B cells. It is well known that Th2 cytokines such as IL-4 and IL-5 are essential for IgE production and IgE-mediated allergic responses because of their function in recruiting B cells, mast cells, and eosinophils involved in allergic inflammatory reactions (14,15). In contrast, the Th1 cytokine IFN- $\gamma$  has inhibitory effects on both IgE production (13) and Th2 differentiation (16). The balance between Th1 and Th2 cytokines is therefore considered crucial for IgE production. In contrast, MDZ significantly increased total IgG production (data not shown). Since IFN- $\gamma$  production was enhanced by MDZ, it is thought that IgG production was also derived from MDZ. The results of *in vivo* cytokine production analyses indicated that i.p. administration of MDZ has an immunomodulatory effect that results in a Th1-skewed cytokine response. Therefore, it is possible that MDZ modulates the balance between Th1 and Th2 responses through induction of Th1 responses



**Fig. 5** Serum levels of IFN- $\gamma$  and IL-4 cytokines.

BALB/c mice were pre-administered MDZ or PBS and then immunized with OVA plus nCT, as described in the legend to Fig. 1. Plasma samples were collected on day 21 and the levels of IFN- $\gamma$  (A) and IL-4 (B) were determined using a Quantikine IFN- $\gamma$  and IL-4 immunoassay kit (R&D Systems, Minneapolis, MN). Values represent mean values  $\pm$  SD ( $n = 5$ ). \* $P < 0.05$  compared with control (OVA plus nCT + PBS). \*\* $P < 0.005$  compared with control (OVA plus nCT + PBS).

in systemic immunity, thereby leading to suppression of IgE production. This is the first demonstration that MDZ administered via the i.p. route can lower the level of Ag-specific IgE in serum.

MDZ significantly inhibited CSR to IgE through inhibition of  $\epsilon$ GLT and  $\epsilon$ CT expression. In contrast, the expression of Id2 was increased by MDZ. Id2 suppresses IgE CSR by inhibiting the  $\epsilon$ GLT-inducing activities of E2A and Pax5 (17,18). It has been shown that TGF- $\beta$  inhibits IgE CSR through the induction of Id2 (5). Therefore, MDZ may increase Id2 expression via the secretion of TGF- $\beta$ , since MDZ was able to enhance TGF- $\beta$  production in serum (data not shown).

Naïve CT is a complex molecule with multiple antigenic and metabolic effects that binds to GM1 gangliosides on the surface of target cells, eventually resulting in increased adenylate cyclase activity and thereby increasing the intracellular concentration of cAMP, resulting in secretion of  $H_2O$ ,  $Na^+$ ,  $K^+$ , and  $HCO_3^-$  into the intestinal lumen, in addition to its antigenic effects (19). As a mucosal adjuvant, CT given via the oral route induces Th2 cells that secrete high levels of IL-4 (20,21). The IL-4 cytokine provides a helper signal for the induction of IgE Abs that may cause anaphylactic reactions (22,23). Indeed, oral and parenteral administration of CT has been reported to induce Th2 responses, such as IgE Ab production, in rodents and humans (24-26). CT induces maturation of human dendritic cells (DC) and licenses them for Th2 priming (27). Since allergic sensitization occurs when antigen-presenting cells such

as DC in the lymphatic system signal T cells, which then interact with B cells to induce IgE production, MDZ may inhibit the adjuvant effect of CT on DC.

In summary, although the mechanism(s) by which MDZ alters the Th1/Th2 balance are yet to be elucidated at the cellular and molecular levels, this is the first demonstration that i.p. administration of MDZ can lower serum IgE levels *in vivo*. These findings suggest that MDZ may be an important modulator of allergic responses through its ability to downregulate IgE production.

### Acknowledgments

We would like to thank Dr. Tomomi Hashizume-Takizawa (Department of Oral Immunology, Nihon Univ. Sch. of Dent. at Matsudo), Dr. Bhawal Kumar Ujjal (Department of Biochemistry and Molecular Biology, Nihon Univ. Sch. of Dent. at Matsudo) and Dr. Michiharu Shimosaka (Department of Anesthesiology, Nihon Univ. Sch. of Dent. at Matsudo) for their helpful suggestions. We would also like to thank Professor Koh Shibutani (Department of Anesthesiology, Nihon Univ. Sch. of Dent. at Matsudo) and Professor Masafumi Yamamoto (Department of Oral Immunology, Nihon Univ. Sch. of Dent. at Matsudo) for their critical reading of this paper and their helpful comments. This work was supported by a grant-in-aid for Challenging Exploratory Research (25670676) from the Japan Society for the Promotion of Science.

### References

- Mertes PM, Laxenaire MC (2004) Allergy and anaphylaxis in anaesthesia. *Minerva Anesthesiol* 70, 285-291.
- Fahy JV (2006) Anti-IgE: lessons learned from effects on airway inflammation and asthma exacerbation. *J Allergy Clin Immunol* 117, 1230-1232.
- Snapper CM, Finkelman FD, Paul WE (1988) Differential regulation of IgG1 and IgE synthesis by interleukin 4. *J Exp Med* 167, 183-196.
- Mao CS, Stavnezer J (2001) Differential regulation of mouse germline Ig gamma 1 and epsilon promoters by IL-4 and CD40. *J Immunol* 167, 1522-1534.
- Sugai M, Gonda H, Kusunoki T, Katakai T, Yokota Y, Shimizu A (2003) Essential role of Id2 in negative regulation of IgE class switching. *Nat Immunol* 4, 25-30.
- Geha RS, Jabara HH, Brodeur SR (2003) The regulation of immunoglobulin E class-switch recombination. *Nat Rev Immunol* 3, 721-732.
- Casellas P, Galiegue S, Basile AS (2002) Peripheral benzodiazepine receptors and mitochondrial function. *Neurochem Int* 40, 475-486.
- Weissman BA, Raveh L (2003) Peripheral benzodiazepine receptors: on mice and human brain imaging. *J Neurochem* 84, 432-437.
- Miyawaki T, Sogawa N, Maeda S, Kohjitani A, Shimada M (2001) Effect of midazolam on interleukin-6 mRNA expression in human peripheral blood mononuclear cells in the absence of lipopolysaccharide. *Cytokine* 15, 320-327.
- Wilms H, Claasen J, Röhl C, Sievers J, Deuschl G, Lucius R (2003) Involvement of benzodiazepine receptors in neuroinflammatory and neurodegenerative diseases: evidence from activated microglial cells in vitro. *Neurobiol Dis* 14, 417-424.
- Rivera R, Segnini M, Baltodano A, Pérez V (1993) Midazolam in the treatment of status epilepticus in children. *Crit Care Med* 21, 991-994.
- Koul R, Chacko A, Javed H, Al Riyami K (2002) Eight-year study of childhood status epilepticus: midazolam infusion in management and outcome. *J Child Neurol* 17, 907-909.
- Pène J, Rousset F, Brière F, Chrétien I, Paliard X, Banchereau J et al. (1988) IgE production by normal human B cells induced by alloreactive T cell clones is mediated by IL-4 and suppressed by IFN-gamma. *J Immunol* 141, 1218-1224.
- Djukanović R, Wilson JW, Britten KM, Wilson SJ, Walls AF, Roche WR et al. (1990) Quantitation of mast cells and eosinophils in the bronchial mucosa of symptomatic atopic asthmatics and healthy control subjects using immunohistochemistry. *Am Rev Respir Dis* 142, 863-871.
- Hamelmann E, Gelfand EW (1999) Role of IL-5 in the development of allergen-induced airway hyperresponsiveness. *Int Arch Allergy Immunol* 120, 8-16.
- Gajewski TF, Fitch FW (1988) Anti-proliferative effect of IFN-gamma in immune regulation. I. IFN-gamma inhibits the proliferation of Th2 but not Th1 murine helper T lymphocyte clones. *J Immunol* 140, 4245-4252.
- Gonda H, Sugai M, Nambu Y, Katakai T, Agata Y, Mori KJ et al. (2003) The balance between Pax5 and Id2 activities is the key to AID gene expression. *J Exp Med* 198, 1427-1437.
- Sugai M, Gonda H, Nambu Y, Yokota Y, Shimizu A (2004) Role of Id proteins in B lymphocyte activation: new insights from knockout mouse studies. *J Mol Med (Berl)* 82, 592-599.
- Merritt EA, Sarfaty S, van den Akker F, L'Hoir C, Martial JA, Hol WG (1994) Crystal structure of cholera toxin B-pentamer bound to receptor GM1 pentasaccharide. *Protein Sci* 3, 166-175.
- Xu-Amano J, Kiyono H, Jackson RJ, Staats HF, Fujihashi K, Burrows PD et al. (1993) Helper T cell subsets for immunoglobulin A responses: oral immunization with tetanus toxoid and cholera toxin as adjuvant selectively induces Th2 cells in mucosa associated tissues. *J Exp Med* 178, 1309-1320.
- Yamamoto M, McGhee JR, Hagiwara Y, Otake S, Kiyono H (2001) Genetically manipulated bacterial toxin as a new generation mucosal adjuvant. *Scand J Immunol* 53, 211-217.
- Kelso JM, Jones RT, Yunginger JW (1993) Anaphylaxis to measles, mumps, and rubella vaccine mediated by IgE to gelatin. *J Allergy Clin Immunol* 91, 867-872.
- Marinaro M, Staats HF, Hiroi T, Jackson RJ, Coste M, Boyaka PN et al. (1995) Mucosal adjuvant effect of cholera toxin in mice results from induction of T helper 2 (Th2) cells and IL-4. *J Immunol* 155, 4621-4629.
- Tamura S, Shoji Y, Hasiguchi K, Aizawa C, Kurata T (1994) Effects of cholera toxin adjuvant on IgE antibody response to orally or nasally administered ovalbumin. *Vaccine* 12, 1238-1240.

25. Snider DP, Marshall JS, Perdue MH, Liang H (1994) Production of IgE antibody and allergic sensitization of intestinal and peripheral tissues after oral immunization with protein Ag and cholera toxin. *J Immunol* 153, 647-657.
26. Qadri F, Asaduzzaman M, Wennerås C, Mohi G, Albert MJ, Abdus Salam M et al. (2000) Enterotoxin-specific immunoglobulin E responses in humans after infection or vaccination with diarrhea-causing enteropathogens. *Infect Immun* 68, 6077-6081.
27. Gagliardi MC, Sallusto F, Marinaro M, Langenkamp A, Lanzavecchia A, De Magistris MT (2000) Cholera toxin induces maturation of human dendritic cells and licenses them for Th2 priming. *Eur J Immunol* 30, 2394-2403.